



**National Registry of Emergency Medical Technicians  
Advanced Level Psychomotor Examination**

**SUPRAGLOTTIC AIRWAY DEVICE**

Candidate: \_\_\_\_\_ Examiner: \_\_\_\_\_  
 Date: \_\_\_\_\_ Signature: \_\_\_\_\_  
 Device: \_\_\_\_\_

NOTE: If candidate elects to initially ventilate with BVM attached to reservoir and oxygen, full credit must be awarded for steps denoted by “\*\*\*” so long as first ventilation is delivered within 30 seconds.

Actual Time Started: _____	Possible Points	Points Awarded
Takes or verbalizes appropriate PPE precautions	1	
Opens the airway manually	1	
Elevates tongue, inserts simple adjunct [oropharyngeal or nasopharyngeal airway]	1	
<b>NOTE: Examiner now informs candidate no gag reflex is present and patient accepts adjunct</b>		
**Ventilates patient immediately with bag-valve-mask device unattached to oxygen	1	
**Ventilates patient with room air	1	
<b>NOTE: Examiner now informs candidate that ventilation is being performed without difficulty and that pulse oximetry indicates the patient's blood oxygen saturation is 85%</b>		
Attaches oxygen reservoir to bag-valve-mask device and connects to high-flow oxygen regulator [12 – 15 L/minute]	1	
Ventilates patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds) with appropriate volumes	1	
<b>NOTE: After 30 seconds, examiner auscultates and reports breath sounds are present and equal bilaterally and medical direction has ordered insertion of a supraglottic airway. The examiner must now take over ventilation.</b>		
Checks/prepares supraglottic airway device	1	
Lubricates distal tip of the device [may be verbalized]	1	
<b>NOTE: Examiner to remove OPA and move out of the way when candidate is prepared to insert device.</b>		
Positions head properly	1	
Performs a tongue-jaw lift	1	
Inserts device to proper depth	1	
Properly secures device per manufacturer's instructions	1	
Ventilates patient and confirms proper ventilation by auscultation bilaterally over lungs and over the epigastrium	1	
Adjusts ventilation as necessary	1	
Verifies proper tube placement by secondary confirmation such as capnography, capnometry, EDD or colorimetric device	1	
<b>NOTE: The examiner must now ask the candidate, "How would you know if you are delivering appropriate volumes with each ventilation?"</b>		
Secures device or confirms that the device remains properly secured	1	
Ventilates patient at proper rate and volume while observing capnography/capnometry and pulse oximeter	1	
<b>Actual Time Ended: _____</b>	<b>TOTAL</b>	<b>18</b>

**Critical Criteria**

- \_\_\_ Failure to initiate ventilations within 30 seconds after using appropriate PPE or interrupts ventilations for greater than 30 seconds at any time
- \_\_\_ Failure to take or verbalize appropriate PPE precautions
- \_\_\_ Failure to voice and ultimately provide high oxygen concentration [at least 85%]
- \_\_\_ Failure to ventilate the patient at a rate of 10 – 12/minute (1 ventilation every 5 – 6 seconds)
- \_\_\_ Failure to provide adequate volumes per breath [maximum 2 errors/minute permissible]
- \_\_\_ Failure to pre-oxygenate patient prior to insertion of the supraglottic airway device
- \_\_\_ Failure to insert the supraglottic airway device at a proper depth or location within 3 attempts
- \_\_\_ Failure to properly secure device per manufacturer's instructions
- \_\_\_ Failure to confirm that patient is being ventilated properly by auscultation bilaterally over lungs and over the epigastrium
- \_\_\_ Insertion or use of any adjunct in a manner dangerous to the patient
- \_\_\_ Failure to manage the patient as a competent EMT
- \_\_\_ Exhibits unacceptable affect with patient or other personnel
- \_\_\_ Uses or orders a dangerous or inappropriate intervention

**You must factually document your rationale for checking any of the above critical items on the reverse side of this form.**